



# Demographic Study of Pattern and Organisms Causing Lung Infection (Pneumonia) In Neurosurgical ICU Setup after Undergoing Neurosurgery

Upadhyay, P. K.<sup>1\*</sup>, Tiwary, G.<sup>2</sup>, Upadhyay, K.<sup>3</sup> & Kritika, U.<sup>3</sup>

<sup>1</sup>Head of the Department, Institute of Human Behavior and Allied Sciences Hospital, University of Delhi, New Delhi, India

<sup>2</sup>Department of Director Laboratory Services and head of Pathology, House of Diagnostic, Delhi, India

<sup>3</sup>MBBS fellow (JSSMC & MAMC)

\*Correspondence to: Dr. Upadhyay, P. K., Head of the Department, Institute of Human Behavior and Allied Sciences Hospital, University of Delhi, New Delhi, India.

## Copyright

© 2022 Dr. Upadhyay, P. K., *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: 24 December 2021

Published: 06 January 2022

**Keywords:** *Pneumonia; Ventilator; Consolidation; Hospital Acquired Infection; Hospital Associated Infection; Kerby- Bauer Method*

## Abstract

Ventilators induced pneumonia are usually associated with some predisposing risk factors. Predisposing factor which were taken into account during this study were prior antacids, prior aspiration, alcoholism, smoking, prior hospital stay and underlying diseases which includes chronic lung disease, diabetes mellitus, neoplasia and heart insufficiency, antibiotic therapy. 50 such case undergone neurosurgery were studied and detailed analysis is presented.

## Introduction

Pneumonia is the most important complication in patient on assisted ventilation. It is also the most

frequent hospital associated infection leading to death. The normal defence mechanism in these patients are bypassed and respiratory tract is led open to colonisation. This is undesirable since at that

situation bacteria are placed advantageously to invade lung parenchyma.

In addition, the result of culture of lung aspirate, which are contaminated with tracheal organism, are made more difficult to interpret.

The bacteria involved have been shown to originate from different sources e.g. Respiratory equipment, nasogastric feeding, other patient and endogenous sources as faeces and other colonised sites in patient. Bacteria from these sources have been found on hands of staff and other vehicles of spread, but route of entry in trachea is less clear. There are two potential common pathways by which bacteria may enter in to trachea -

- 1) Through the airway of endotracheal tube or tracheostomy tube via inhaled air during suction.
- 2) Via contaminated oropharyngeal or gastric secretion passing between endotracheal tube and trachea.

Bacteria passes through a tracheostomy tube wound can also occur, but is less important because wound receives protection from infection.

The air delivered to patient may carry bacteria entrapped from a contaminated part of respiratory equipment or from air of unit itself. However much emphasis is placed on preventing bacterial contamination of respiratory equipment and ventilators are generally decontaminated after use by efficient processes. Humidifiers are generally filled with sterile water and are stored dry between patients. Respiratory tubing's are changed frequently.

Less work has been done on bacterial contamination of air of i.c.u., but common experience and available evidence tend to indicate that it is of little importance. In any case, in cross infection episode, the pattern of spread of tracheal isolates between patients is typical of contact spread. Contamination of inspired air from retrograde movements of

tracheal secretion in to respiratory line could occur after tracheal colonisation.

## Material and Method

Fifty patients in the neuro-surgery i c u of our hospital receiving mechanical intubation were studied. Patients were randomly selected.

Patients with short term (<48 hrs.), and long term ventilation therapy (>48 hrs) were included. Each patient was ventilated via an endotracheal or tracheostomy tube connected to one of the following mechanical ventilators.

## Data Collection

Daily patient temperatures, laboratory results, underlying disease and drug therapy was recorded prospectively on standardized forms.

The diagnosis of ventilation associated pneumonia required four of the following five criteria developing not less than 48 hours after the start of mechanical ventilation:

- 1) A new and persistent infiltrate in a chest x-ray.
- 2) Purulent tracheal secretion
- 3) A significant pathogen isolated from cultures of tracheal aspirate
- 4) Body temperature above 38
- 5) Peripheral leukocyte count of more than 10,000/cumm and

## Physical Signs of Pneumonia

Cough, fever, sputum discharge, breathlessness, fall in oxygen saturation in various combinations. Along with these criteria, a positive chest x-ray must always be present.

Coma was defined as score of 8 by the Glasgow coma score. Most of the patient were between 5 and 7 of g.c.s. The acute physiological score quantities abnormalities in the patients vital signs, mental status, and organ function on the basis of the most abnormal values and physical signs with 24 hours of admission to i.c.u.

Medication such as antibiotics, steroids, antacids, vasopressors and pancuronium bromide (pavlon) and bronchodilators were recorded. Each medication was considered a potential risk factor for pneumonia if given when patient was receiving mechanical ventilation and prior to pneumonia. Personal history of the patient regarding alcoholism, smoking were recorded which also included in predisposing factors.

Other predisposing factors are also recorded which included prior aspiration, prior hospital stay, chronic lung disease, diabetes, neoplasm and heart insufficiency.

### Microbiological Sampling

- 1) Reservoir samples were collected after 24hrs of ventilation and promptly cultured for bacteria and fungi. All plates were incubated at 37°C for 48 hrs.
- 2) trachea-bronchial secretions - culture of tracheal secretions were collected by suction directly from the lower respiratory tract via the tracheostomy or endo tracheal tube organisms were identified according to routine microbiological methods. No cultures were made for anaerobic bacteria, legionella, mycoplasma, or viruses.
- 3) Other cultures: - nasal and throat swab were collected to identify organism being carried at these sites.
- 4) Blood and urine culture were also done in the entire patient to exclude any previous infection in the patients.

Antibiotic sensitivity of different organism was done by the Kerby- Bauer method. Sensitivity test were done for the following antibiotics - ampicillin, cephaazolin, chloram phenicol, carbenicillin, kanamycin, cotrimoxazole, tetracycline, cloxaacillin, erythromycin, lincomycin, cefotaxime, ceftazidime, streptomycine, colistin, gentamycin, polymixin -b. Strains with similar antibiotic sensitivity pattern up to a difference of two were considered to be the same.

Procedure for decontamination of ventilator apparatus:

- 1) Ventilator tubing changed every 48 hrs and decontaminated by ethylene oxide gas.
- 2) Reservoir fluid was changed daily
- 3) A new suction catheter used for every suction procedure.

Patients included in the study were suffering from different neurosurgical problems which included the broad subgroups as supratentorial mass lesion, vascular pathology, mainly aneurysm, and a.v.m., posterior fossa masses include both midline and c.p. angle brain stem masses, hydrocephalus - congenital and post meningitic, head injuries, and high cervical cord compression. Studied patient belongs to different age groups.

### Age Variations

Youngest patient was 4 years old and oldest was 78 years old. Most of the patients were between 3rd and 5<sup>th</sup> decades.

Demographic characteristic in the 50 studied patient are shown in table 1, 2 and 3;

**Table 1:** Age of the patients

Age in years	No. of patient
1 -10	7(14%)
11-20	4
21-30	7
31-40	12(24%)
41-50	7
Above 50	13(26%)
Total	50

**Table 2**

Sex	No. of patients
Male	38(76%)
Female	12(24%)
Total	50

**Table 3:** Clinical diagnosis at admission

1) Supratentorial mass lesion	19(38%)
2) Infra tentorial mass lesion	10(20%)
3) Hydrocephalus	6
4) Vascular lesion	8
5) High cervical cord compression	3
6) Head injury	4
Total	50

Characteristic of the 50 patients on admission to the i.c.u. are shown in table 4.all these patients were between 5 and 7 of g.c.s.

**Table 4:** Characteristics of 50 patients on Mechanical, ventilation

Characteristics	No. of patients
Prior aspiration	22 (44%)
Prior antacids	26 (52%)
Alcoholism	4 (8%)
History of smoking	20 (40%)

<b>Prior hospital stay</b>	
1 day	4 (8%)
2-7 days	8 (16%)
More than 7 days	38 (76%)
Antibiotics	37 (74%)
Glasgow coma scale (between 5-7)	50 (100%)
Underlying disease (chronic lung diseases), diabetes, neoplasia, heart insufficiency	23 (46%)

Most of the patients studied were already present in the hospital for more than two days. Out of which 22 patients (44%) had history of prior aspiration, 26 patients (52%) of prior antacids, 4 (8%) patient had history of alcoholism and 20(40%) patient has history of smoking. Hospital stay was 1 day in 4 (8%) patient, 2-7 days in 8 (16%) patients, more than 7 days in 38 (76%) patient, 23 (46%) had underlying

disease in form of chronic lung disease, neoplasia or heart disease. Thirty seven patients were already on antibiotics.

Out of the 50 patient included in study, 23 develops pneumonia. The probable predisposing factors to the development of pneumonia in these patient are given in table 5.

**Table 5**

<b>Predisposing factors</b>	<b>No. of patients</b>	<b>Pneumonia</b>
❖ Prior aspiration	22 (44%)	15 (68.1%)
❖ Prior antacid	26 (52%)	11 (42.3%)
❖ Alcoholism	4 (8%)	2(50%)
❖ Smoking	20 (40%)	9 (45%)
❖ Prior hospital stay		
1 day	4 (8%)	1 (25%)
2-7 days	8 (16%)	5 (62.5%)
More than 7 days	38 (76%)	17 (44.7%)
❖ Antibiotics	37 (74%)	23 (66%)
❖ Underlying diseases	23 (46%)	8 (30.4)%
❖ Duration of ventilation		
Less than 48 hours	6	1 (28%)
More than 48 hours	44	22 (50%)
<b>Total:</b>	<b>50</b>	<b>23 (46%)</b>
❖ <b>Significant</b>		

Five independent variables were significantly associated with ventilator as associated pneumonia. Histories of prior aspiration were found in 22 (44%) patients out of whom 15 patients (68.1%) associated

with pneumonia. Histories of alcoholism were present in 4 patients (8%) of which 2 patents (50%) associated with pneumonia. Other significant factors were prior hospital stay between 2-7 days

administration of antibiotics and more than 48 hours duration on ventilator.

**Bacteriological Study**

The organisms isolated from various sites in the 50 study patients are shown in table 6.

*Table 6: Organism isolated from various site in 50 patients on mechanical ventilation*

Organisms	Tracheal	Reservoir	Urine*	Nosc/throat**	Total
P. Aeruginosa	9	6	4	1	20
K. Pneumoniae	26	1	9	2	38
E. Coli	2	1	6	2	12
Staph aureus	5	-	-	2	7
Proteus sp.	1	-	-	-	1

The gram negative bacilli were the most frequent group of organisms isolated from the different sites. Klebsiella pneumonia was most common, next was pseudomonas aeruginosa followed by escherichia coli.

\* Similar organisms in urine and tracheal aspirate were found in 10 patients. In 9 patients organisms were different.

\* Similar organisms in nose/throat, and tracheal aspirate were found in 2 patients. In 5 patients organisms were different.

The various organisms isolated from the tracheal

aspirate of the 23 cases, which developed pneumonia is given in table 7.

Gram negative bacteria were the predominant organisms being isolated from 82% of tracheal aspirates of patients who developed ventilator associated pneumonia. In many of the aspirates more than one organism were isolated. The most common organism isolated were klebsiella species (60.1%), pseudomonas aeruginosa (39.1%), proteus species (8.9%).

Pneumonia developed in the majority of cases (80%) after 7 days of ventilator therapy.

*Table 7: Reservoir fluid colonized with bacteria*

Organisms	No. of patients
P. Aeruginosa	6
Klebsiella pneumonia	1
E. Coli	1
Aerobic spore bearers	1
Acinetobactor sp.	1
Total:	10 (20%)

Six of these 10 patients acquired ventilator associated pneumonia. In 6 patients, the bacteria isolated from the reservoir sample (one klebsiella

pneumoniae, 5 pseudomonas aeruginosa) were found to be the same organisms as cultured from trachea secretions. All the six patient in whom the

reservoir fluid and tracheotomy or intubation aspirate culture were same, developed pneumonia; whereas 3 of the remaining 4 patients where there was no correlation between reservoir fluid and tracheotomy or aspirate bacteriology developed pneumonia. In these patients tracheal secretions were probably the source of reservoir colonization while

in the four patients where the bacteria isolated from the reservoir samples (one E. coli, one pseudomonas sp., one aerobic spore bearing organism and one acinetobactor sp.) did not correspond with those recovered from tracheal aspirates, tracheal secretions were probably not the source of contamination of reservoir (table 8).

**Table 8:** Bacteriology of tracheal aspirate and reservoir Fluid in 9 cases of pneumonia in whom The ventilator humidifier was contaminated

Organisms grown in Reservoir fluid	Organisms grown In tracheostomy Or endotracheal Aspirate	No. Of patients	Pneumonia
P. Aeruginosa	Pseudomonas Aeruginosa	5	5
Klebsiella pneumonia	Klebsiella pneumoniae	1	1
E. Coli	Klebsiella pneumoniae	1	1
Aerobic spore Bearing bacilli	Klebsiella pneumoniae	1	-
Pseudomonas sp.	Proteus Mirabilis	1	1
Acinetobactor sp.	Klebsiella pneumonia	1	1
<b>Total:</b>		<b>10</b>	<b>9</b>

Most of the patients, whose prior hospital stay were between 2-7 days or more than 7 days showed organisms in the tracheostomy/intubation aspirate or nasal/throat swab or urine. Mechanical ventilator induced pneumonia by nosocomial organisms were more common in patients with prolonged hospital stay.

Duration of mechanical ventilation is also important. Patients, who were electrically ventilated for less than 48 hours, did not show any evidence of infection barring a few patients. Mostly infection and mechanical ventilators induced pneumonia seen in patients who were ventilated for more than 48 hours.

**Table 9:** Corelation between ventilators associated pneumonia and culture positive tracheal aspirate, reservoir Fluid, blood and urine culture

	Ventilator Associated Pneumonia	No ventilation Associated Pneumonia	Total
Total no. Of patients with ventilators	23	27	50
Tracheal aspirate	23	19	42
Reservoir fluid	9	1	10
Urine	13	6	19
Blood	2	-	2

## Discussion

Previous investigators have identified several risk factors for the development nosocomial pneumonia [1]. Although most patients who develop hospital acquired pneumonia do not have artificial airways, patients who receive mechanical ventilator via an orotracheal or nasotrachea tube have a substantially increased risk of lower respiratory tract infection [2]. Both orotracheal and nasotracheal tubes bypass natural defences, permit leakage of bacteria and secretions around the cuff into the trachea and reduce bacterial clearance from the trachea. The precise diagnosis of pneumonia in patients receiving mechanical ventilation is often difficult. Most of the patients have serious disease, increased oropharyngeal, colonization by hospital flora and numerous reactive elevated body temperature or leucocytosis [3-6]. Purulent sputum may occur because of intubation or leakage of oropharyngeal secretions around the artificial airway. Furthermore, chest roentgenogram changes consistent with pneumonia may be caused by pulmonary oedema, pulmonary infraction, atelectasis are used as established criteria for the diagnosis of pneumonia, the clinical assessment of at least two independent investigators was considered necessary before diagnosing pneumonia, although we can include the possibility that pneumonias are incorrectly diagnosed, it is unlikely any true cases of pneumonia would have been missed by our criteria. The overall incidence of ventilator induced pneumonia in our study was 46%. Other studies have found pneumonia rates ranging from 21% to 66% [2,7,8].

Ventilators induced pneumonias are usually associated with some predisposing risk factors. Predisposing factor which were taken into account during this study were prior antacids, prior aspiration, alcoholism, smoking, prior hospital stay and underlying diseases which includes chronic lung disease, diabetes mellitus, neoplasia and heart insufficiency,

antibiotic therapy, duration by Craven DE *et al* (1986), F. Daschner *et al* (1987) [7,8]. Out of all the factors, mechanical ventilator associated pneumonia was more common in patients having prior aspiration (68.1%), alcoholism (50%), smoking (45%) craven *et al* (1986) [7] considered the raised intracranial pressure, cimetidine and antacids, fall-winter season as a significant variables for ventilator associated pneumonia. Their results may reflect severely ill patient at high risk for pneumonia rather than a direct causal relationship with pneumonia. When intracranial pressure increases, chest physiotherapy and frequent suctioning of pulmonary secretions are often avoided, special precautions are taken in neurosurgical patients, prevent repositioning of patient after surgery for CV junction or high cervical cord.

Gastrointestinal bleeding in critically ill patients can be reduced by the use of antacids and cimetidine [9]. Decreased gastric acidity increased gastric and tracheal colonization particularly when a gastric tube is in place [4,9,10]. In our study, 42.3% patients given antacids or ranitidine developed pneumonia, which is reported to be 37% by craven *et al* 1986. Ranitidine causes overgrowth of bacteria in the stomach [4,10], with tracheal colonization or may allow the host inflammatory and immune response decrease in the lung [11].

This study does not support the previous data suggesting that mechanical ventilators circuits should be changed every 48 hour rather than every 24 hour as recommended by the centres for disease control 3, craven *et al* 1986 [7]. It was emphasized that ventilators without mainstream reservoir nebulizer impose no greater risk to a patient that imposed by breathing room air. Most mechanical ventilators have air that is moistened by cascade humidifier rather than by a reservoir nebulizer. The significant increase rate of pneumonia in patients having 24 hour ventilator circuit changes may have resulted

from increased from increased manipulation of the patient, endotracheal tube or ventilator tube as suggested by Craven *et al* (1986) [7], but this was not seen in this study. They suggested the cause to be a probable contamination of the patients' airway by contaminated tubing condensate [12] or increased leakage of bacteria around the endotracheal tube into the trachea. Similarly the increased manipulation of other devices such as intravenous or bladder catheters may also increase nosocomial infection [13]. In the present study, similar organisms were seen in the urine and tracheal aspirate in ten patients. This could have been due to transmission through nurses hands while manipulation urinary catheter following or preceding the suction procedures. Gram negative bacilli were frequently isolated from the sputum of patients with pneumonia as also seen by LA Force *et al* (1981) [14], Dashner *et al* 1988) [8], Craven *et al* (1986) [7], and Bryan *et al* (1984) [6]. Increased colonization of the pharynx with gram negative bacilli may be due to the severity of the patient's underlying disease or the use of drugs such as ranitidine. In the present study, the use of antibiotics was as associated with higher incidence of pneumonia. Antibiotics not only alter colonization in the oropharynx and gastrointestinal tract but also may have deleterious effects on the host immune response to infection (Hanser WE *et al* 1982) [15].

However, the relationship between antibiotic use and nosocomial pneumonia is difficult to assess because of the wide spectrum of agents used alone or in combination in the patient's population.

Use of respiratory filters [16,17] in the inspiratory and expiratory lines on the patient's side of the ventilator are used to prevent colonization of the ventilatory circuit by bacteria. Contamination of the lines between the patients and the filters was prevented by autoclaving the respiratory tubing every 24 hours and by filling humidifier reservoir with 0.02% chlorhexidine. A sterile technique used for

tracheal suction, but rate of recovery of pseudomonas from tracheal aspirates was similar to that found before these measures were undertaken [16,17]. Suction of the tracheobronchial tree may lead to contamination of the trachea into the trachea during insertion of catheter.

In the present study, ventilatory circuit was decontaminated every 24 hour and reservoir fluid changed daily. However in 10 case contamination of reservoir fluid occurred. The contamination organism in 6 cases was the same as that isolated from tracheal secretions. Tracheal secretions were probably the source of reservoir contamination in these cases. Contamination could have taken place via contaminated tubing condensate or during manipulation.

Organisms are known to enter the mouth by various means, particularly in foods and fluids and during mouth washing (Johnson *et al* 1969) [3]. Mouth washing is always regarded as a sterile procedure and has been carried out with tap waters and cleaning pots contaminated with bacteria. Swallowed mouth secretions and nasogastric feed will introduce hospital acquired organisms into the stomach, where of antacids and H blockers are used, they will multiply. Consequently gastric regurgitation is another source of bacteria which may enter the respiratory passages by final common pathway of leakage around the endotracheal tube.

The recovery of organisms in the pharyngeal and tracheal secretions could also be explained by their deposition from contaminated hands in the mouth during oral hygiene and in the endotracheal tube during tracheal suction.

The danger of this route of spread can be reduced in various ways:

1. By preventing the entry of organisms into the mouth and stomach by providing sterile feeds and regular oral disinfection.

2. By avoiding the accumulation of secretions above the endotracheal tube cuff by nursing the patient in a head down position and keep the nasopharynx below trachea.

3. By removing such secretions via a small suction tube attached along the convex border of an endotracheal tube with its tip at the proximal end of the cuff.

## Conclusions

Ventilators induced pneumonias are usually associated with some predisposing risk factors. Predisposing factor which were taken into account during this study were prior antacids, prior aspiration, alcoholism, smoking, prior hospital stay and underlying diseases which includes chronic lung disease, diabetes mellitus, neoplasia and heart insufficiency, antibiotic therapy.

Out of all the factors, mechanical ventilator associated pneumonia was more common in patients having prior aspiration (68.1%), alcoholism (50%), smoking (45%) craven *et al* (1986) considered the raised intracranial pressure, cimetidine and antacids, fall-winter season as a significant variables for ventilator associated pneumonia.

Swallowed mouth secretions and nasogastric feed will introduce hospital acquired organisms into the stomach. In Patients given antacids or histamine-2 blocker developed pneumonia more. Ventilators without mainstream reservoir nebulizer impose no greater risk to a patient for pneumonia. Our study does not support the previous data suggesting that mechanical ventilators circuits should be changed every 48 hour rather than every 24 hour as recommended by the centres for disease control. The danger of this route of spread can be reduced in various ways:

4. By preventing the entry of organisms into the mouth and stomach by providing sterile feeds and regular oral disinfection.

5. By avoiding the accumulation of secretions above the endotracheal tube cuff by nursing the patient in a head down position and keep the nasopharynx below trachea.

6. By removing such secretions via a small suction tube attached along the convex border of an endotracheal tube with its tip at the proximal end of the cuff.

## Results

Pneumonia developed in the majority of cases (80%) after 7 days of ventilator therapy.

K. Pneumoniae was found in 38 cases P. Aeruginosa was found in 20 cases, E. coli in 12 cases, and Staph aureus in 7 cases and proteus in 1 case (Details of places involved are given in table 6).

Reservoir fluid was found contaminated in 10 cases (20%).

Total of fifty cases 23 case were found to be having ventilator associated pneumonia, while 27 were having non ventilator associated pneumonia. Ventilator associated pneumonia have 100% tracheal aspirate positive while only 19 case in non-ventilator associated. Likewise 9 out of 23 had reservoir fluid positive culture while only one in non-ventilator induced pneumonia (detail see table 9).

## Bibliography

1. Hooton, T. M., Haley, R. W., Culver, D. H., *et al.* (1981). The joint association of multiple risk factors with the occurrence of nosocomial infections. *Am J Med.*, 70(4), 960-970.
2. Cross, A. S. & Roup, B. (1981). Role of respiratory assistance devices in endemic nosocomial pneumonia. *Am J Med.*, 70(3), 681-685.

3. Johnshon, W. G., Pierce, A. K. & Sanford, J. P. (1969). Changing pharyngeal bacterial flora of hospitalized patients. Emergence of gram negative bacilli. *New Engl. J Med.*, 281(21), 1137-1140.
4. Atherton, S. T. & White, D. J. (1978). Stomach as a source of bacteria colonising respiratory tract during artificial ventilation. *Lancet*, 2(8097), 968-969.
5. Rose, H. D. & Babcock, J. B. (1975). Colonisation of icu patients with gram negative bacilli. *Amj Epidermoid.*, 101(6), 495-501.
6. Bryan, C. S. & Reynolds, K. L. (1984). Bacteremic nasocomial pneumonia. *Am rev resp. Dis.*, 129(5), 668-671.
7. Craven, D. E., Kunches, L. M., Kilinsky, V., Lichtenberg, D. A., Make, B. J. & Mccobe, W. R. (1986). Risk factors for pneumonia and fatality in patients receiving continuous mechanical ventilation. *Am Rev Resp Dis.*, 133(5), 792-796.
8. Dashner, F., Kappstein, I., Schuster, F., Scholz, R., Bauer, E., Joobens, D. & Just, H. (1988). Influence of disposable and reusable humidifying system on the incidence of ventilator pneumonia. *Journal of Hospital Infections.*, 11(2), 161-168.
9. Hastings, P. R., Skillman, J. J., Bushness, L. S. & Silen, W. (1978). Antacid titration in the prevention of acute gastrointestinal bleeding - A controlled, randomized trial in 100 critically ill patients. *N. Enld. J Med.*, 298, 1041-1045.
10. Moulin, D. U., Hedley-Whyte, J., Paterson, D. G. & Lisbon, A. (1982). Aspiration of gastric bacteria in antacid treated patients - a frequent cause of postoperative colonization of airway. *Lancet*, 1(8266), 242-245.
11. Beer, D. J., Matlof, S. M. & Rocklin, R. E. (1984). The influence of histamine on immune and inflammatory responses. *Adv. Immunol.*, 35, 209-268.
12. Craven, D. E., Goularte, T. A. & Make Barry, J. (1984). Contaminated condensate in mechanical ventilation circuits. A risk factor for nosocomial pneumonia. *Am Rev Respir Dis.*, 129(4), 625-628.
13. Burke, J. P., Garabaldi, R. A. & Brittt, M. R. (1981). Prevention of catheters associated urinary tract infections, efficacy of daily mental care regime. *Am J Med.*, 70, 542-546.
14. Laforce, F. M. (1981). Hospital acquired gram negative rod pneumonia: an overview. *Am J Medicine.*, 70(3), 664-669.
15. Hauser, W. E. & Remington, J. S. (1982). Effect of antibiotics on the immune response. *Am J Med.*, 72(5), 711-716.
16. Lumley, J. (1976). Decontamination of anaesthetic equipment and ventilator. *Brit J Anaesth*, 48(1), 3-8.
17. Holdcraft, A., Lumbley, J. & Gaya, H. (1973). Why disinfect ventilators? *Lancet*, 1(7797), 240-241.